

University of Colorado Denver
Department of Mathematical and Statistical Sciences
Applied Linear Algebra Ph.D. Preliminary Exam
August 10, 2020

Name: _____

Exam Rules:

- This exam is being administered remotely using Zoom. During the exam, you must be logged in to the assigned Zoom meeting with your camera on, and you must be visible in the camera. Your microphone can be muted, but please leave Zoom audio ON so that the proctor can speak to you if needed.
- If you have any questions during the exam, please contact the exam proctor using the zoom chat, or call the proctor at the number given below.

Committee Member Contact Information:

Name	Phone	email	Proctoring times
Stephen Billups	303-877-2861	stephen.billups@ucdenver.edu	10am-noon
Yaning Liu	850-980-5275	yaning.liu@ucdenver.edu	noon-2pm
Julien Langou	720-520-2358	julien.langou@ucdenver.edu	alternate

- If you need to be out of view of the camera for any reason (bathroom breaks, etc.), please let the proctor know by posting a message on the zoom chat.
- If you would like to work with a paper copy of the exam, please print it as soon as you receive it and inform the proctor that you are doing so before the exam begins. If the printer is in another room, let the proctor know.
- You may read the exam as soon as you receive it, but you may not start writing (even your name) until authorized to start writing.
- This is a closed book exam. You may not use any external aids during the exam, such as:
 - communicating with anyone other than the exam proctor (through text messages or emails, for example);
 - consulting the internet, textbooks, notes, solutions of previous exams, etc;
 - using calculators or mathematical software.
- You may use a tablet PC (such as an iPad or Microsoft Surface) to write your solutions. Alternatively, you can write your solutions on paper.
- Please begin each problem on a new page, and write the problem number and page number at the top of each page. (For example, 6-1, 6-2, 6-3 for pages 1, 2 and 3 of problem 6). If you are writing on paper, write only on one side of the paper.
- The exam will end 4 hours after it begins. At the conclusion of the exam, please email a copy of your solutions to the exam proctor. Do not leave until the proctor acknowledges that your solutions have been successfully received.

- Your solutions need to be in a single .pdf file with the pages in the correct order. The .pdf file needs to be of good enough quality for easy grading.
 - If you cannot create a good quality .pdf file quickly, you may instead submit an imperfect scan, or even pictures of your exam, and then take more time to prepare and submit a good quality .pdf version. We will grade the better version but use the first submission to check that nothing was added or changed between versions.
 - Do not submit your scratch work.
- There are 8 total problems. Do all 4 problems in the first part (problems 1 to 4), and pick two problems in the second part (problems 5 to 8). Do not submit more than two solved problems from the second part. If you do, only the first two attempted problems will be graded. Each problem is worth 20 points.
 - Do not submit multiple alternative solutions to any problem; if you do, only the first solution will be graded.
 - Justify your solutions: **cite theorems that you use**, provide counter-examples for disproof, give explanations, and show calculations for numerical problems.
 - If you are asked to prove a theorem, do not merely quote that theorem as your proof; instead, produce an independent proof.
 - Notation: Throughout the exam, \mathbb{R} and \mathbb{C} denote the sets of real and complex numbers, respectively. \mathbb{F} denotes either \mathbb{R} or \mathbb{C} . \mathbb{F}^n and $\mathbb{F}^{n,n}$ are the vector spaces of n -tuples and $n \times n$ matrices, respectively, over the field \mathbb{F} . $\mathcal{L}(V)$ denotes the set of linear operators on the vector space V . T^* is the adjoint of the operator T and λ^* is the complex conjugate of the scalar λ . In an inner product space V , U^\perp denotes the orthogonal complement of the subspace U .
 - If you are confused or stuck on a problem, either ask a question or move on to another problem.

Problem	Points	Score		Problem	Points	Score
1.	20			5.	20	
2.	20			6.	20	
3.	20			7.	20	
4.	20			8.	20	
				Total	120	

Applied Linear Algebra Preliminary Exam Committee:

Stephen Billups (Chair), Julien Langou, Yaning Liu

Part I. Work **all** of problems 1 through 4.

Problem 1.

(a) Let $A = \begin{bmatrix} 2 & -1 & 3 & -1 & 1 & -5 \\ -1 & 4 & 2 & 11 & -3 & 10 \\ 1 & 1 & 3 & 4 & 2 & -7 \end{bmatrix}$. Find bases for the null and column spaces of A .

(b) Let S and T be subspaces of \mathbb{R}^n .

(i) Show that there exist matrices A and B such that $S = \text{null } A$ and $T = \text{null } B$. Given bases for S and T , describe briefly how such matrices A and B could be found.

(ii) Find a matrix C such that $S \cap T = \text{null } C$.

(iii) Describe briefly how to find a basis for $S \cap T$.

Problem 2.

Let V be a finite-dimensional real inner product space. Let $T \in \mathcal{L}(V)$ and let U be any T -invariant subspace of V . Prove or give a counterexample for each of the following statements:

(a) U^\perp is T^* -invariant.

(b) U^\perp is T -invariant.

Problem 3.

For $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$, let $\langle \mathbf{x}, \mathbf{y} \rangle$ be the Hermitian inner product $\sum_j x_j \bar{y}_j$. Let T be an operator on \mathbb{C}^n such that $\langle T\mathbf{x}, T\mathbf{y} \rangle = 0$ if $\langle \mathbf{x}, \mathbf{y} \rangle = 0$. Prove that $T = kS$ for some scalar k and some operator S which is unitary (an isometry).

Problem 4.

Consider the vector space $\mathcal{M}_{n \times n}(\mathbb{R})$ of $n \times n$ real matrices. Consider the linear map $T : \mathcal{M}_{n \times n}(\mathbb{R}) \rightarrow \mathcal{M}_{n \times n}(\mathbb{R})$ given by $T(A) = A^T$ for all $A \in \mathcal{M}_{n \times n}(\mathbb{R})$. Here A^T denotes the transpose of A .

- Find the characteristic polynomial and minimal polynomial of T .
 - Find the Jordan form of T .
-

Part II. Work **two** of problems 5 through 8.

Problem 5.

Let A be an $n \times n$ matrix over \mathbb{C} . Show that

1. there exists an $n \times n$ invertible matrix Q such that

$$Q^{-1}AQ = \begin{bmatrix} \lambda_1 & b_{12} & \cdots & b_{1n} \\ 0 & b_{22} & \cdots & b_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & b_{n2} & \cdots & b_{nn} \end{bmatrix}$$

2. Use mathematical induction to show that any $n \times n$ matrix over \mathbb{C} is similar to an uppertriangular matrix

$$\begin{bmatrix} \lambda_1 & * & \cdots & * \\ 0 & \lambda_2 & \cdots & * \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}$$

Problem 6.

Let V be a finite dimensional inner product space over \mathbb{C} . Let $T : V \rightarrow V$ be a self-adjoint operator on V . Suppose $\mu \in \mathbb{C}, \epsilon > 0$ are given and assume there is a unit vector $\mathbf{x} \in V$ such that

$$\|T(\mathbf{x}) - \mu\mathbf{x}\| \leq \epsilon$$

Show that there is an eigenvalue λ of T such that $|\lambda - \mu| \leq \epsilon$.

Problem 7.

Let V be an inner product space with inner product $\langle \cdot, \cdot \rangle$.

An operator P in $\mathcal{L}(V)$ is said to be a *projection* if and only if $P^2 = P$.

A projection P is said to be an *orthogonal projection* if and only if $\text{Range}(P)$ and $\text{Null}(P)$ are orthogonal subspaces.

1. Let P be a projection. Prove that P is an orthogonal projection if and only if P is diagonalizable in an orthogonal basis.
 2. Let P be a projection. Prove that P is an orthogonal projection if and only if $\|Pv\| \leq \|v\|$ for every $v \in V$.
-

Problem 8.

Recall that $\text{trace}(A^T B)$ defines an inner product on $\mathcal{M}_{n \times n}(\mathbb{R})$ and that the norm associated with this inner product is $\|\cdot\|_{\text{fro}}$, the Frobenius norm of a matrix. Recall that a symmetric matrix S is such that $S = S^T$ and a skew-symmetric matrix N is such that $N = -N^T$.

Let \mathcal{S} be the subspace of symmetric matrices in $\mathcal{M}_{n \times n}(\mathbb{R})$. Let \mathcal{N} the subspace of skew-symmetric matrices in $\mathcal{M}_{n \times n}(\mathbb{R})$.

1. Prove that

$$\mathcal{S} \oplus \mathcal{N} = \mathcal{M}_{n \times n}(\mathbb{R})$$

2. Prove that the subspaces \mathcal{S} and \mathcal{N} are orthogonal with respect to the inner product $\text{trace}(A^T B)$.
3. Let A be in $\mathcal{M}_{n \times n}(\mathbb{R})$. Prove that

$$\min_{S \in \mathcal{S}} \|A - S\|_{\text{fro}} = \frac{1}{2} \|A - A^T\|_{\text{fro}}.$$
