



## Corrigendum

### Wing size, wing shape and sexual dimorphism in eye-span in stalk-eyed flies (Diopsidae)

GAL RIBAK<sup>1</sup>, MARIE L. PITTS<sup>2</sup>, GERALD S. WILKINSON<sup>2</sup> and JOHN G. SWALLOW<sup>1</sup>

<sup>1</sup>Department of Biology, University of South Dakota, Vermillion, SD 57069, USA

<sup>2</sup>Department of Biology, University of Maryland, College Park, MD 20742, USA

The authors recently discovered that 15 out of the 421 flies measured in the above mentioned study were erroneously measured twice at different time points of the data collection process. Therefore the correct sample size for this analysis (without repeated sampling) should have been 406 flies. The 15 flies that were measured twice belonged to one species (*Diasemopsis meigenii*) and included 7 male flies and 8 females. A re-calculation of the entire statistical analysis, this time with the correct sample size confirmed that the inflated sample size for *D. meigenii* had negligible effect on statistical values and did not change any of the results or conclusions reported in the paper.

For the sake of accuracy we provide the corrected averages for *D. meigenii* below and these should be used instead of the ones presented for *D. meigenii* (DM) in figure 4.

The authors regret any inconvenience that might have resulted from this error.

#### Means $\pm$ 1 SD of measurements from 20 males and 20 females of *D. meigenii*

<i>Diasemopsis meigenii</i>	Body mass [mg]	Wing length [mm]	Eye span [mm]	Wing area [mm <sup>2</sup> ]
Males	10.4 $\pm$ 1.39	4.7 $\pm$ 0.20	8.0 $\pm$ 0.42	10.4 $\pm$ 0.70
Female	10.0 $\pm$ 1.84	4.6 $\pm$ 0.40	5.4 $\pm$ 1.93	10.5 $\pm$ 1.65

#### REFERENCE

Ribak G, Pitts ML, Wilkinson GS, Swallow JG. 2009. Wing size, wing shape and sexual dimorphism in eye-span in stalk-eyed flies (Diopsidae). *Biological Journal of the Linnean Society* **98**: 860–871.